

Harmful molecular mechanisms in sepsis

Daniel Rittirsch, Michael A. Flierl and Peter A. Ward

Abstract | Sepsis and sepsis-associated multi-organ failure are major challenges for scientists and clinicians and are a tremendous burden for health-care systems. Despite extensive basic research and clinical studies, the pathophysiology of sepsis is still poorly understood. We are now beginning to understand that sepsis is a heterogeneous, dynamic syndrome caused by imbalances in the 'inflammatory network'. In this Review, we highlight recent insights into the molecular interactions that occur during sepsis and attempt to unravel the nature of the dysregulated immune response during sepsis.

Sepsis

A systemic response to severe infection or tissue damage, leading to a hyperactive and unbalanced network of pro-inflammatory mediators. Vascular permeability, cardiac function and metabolic balance are affected, resulting in tissue necrosis, multi-organ failure and death.

T_H17 cells

(T helper 17 cells). A subset of CD4⁺ T helper cells that produce interleukin-17 (IL-17) and that are thought to be important in inflammatory and autoimmune diseases. Their generation involves IL-23 and IL-21, as well as the transcription factors ROR γ t (retinoic-acid-receptor-related orphan receptor- γ t) and STAT3 (signal transducer and activator of transcription 3).

The clinical manifestations of sepsis were already known to Hippocrates (460–377 BC), who introduced the term 'wound putrefaction'. In addition, the Persian 'father of modern medicine', Ibn Sina (also known as *Avicenna*, AD 980–1037), observed that septicaemia was usually accompanied by fever. However, it was not until the 18th century that Louis Pasteur linked the decay of organic substances to the presence of bacteria and microorganisms, and Ignaz Semmelweis observed the significant effect of hygienic measures on decreasing the mortality of women during childbirth. In 1914, Hugo Schottmüller laid the foundations for a modern definition of sepsis and was the first to describe that the presence of an infection was a fundamental component of the disease. Decades later, the ideas of Lewis Thomas led to a turnaround in the understanding of sepsis by popularizing the theory that "...it is the [host] response ... that makes the disease"¹. This theory resulted in a large number of experimental and clinical studies, which eventually shifted the focus of sepsis research from the infectious agent to the host immune response. Finally, the concept entered into daily clinical practice when Roger Bone and colleagues defined sepsis as a systemic inflammatory response syndrome (SIRS) that can occur during infection².

In the past, sepsis was commonly thought to be caused by overactivation of the innate immune system, and the ensuing pro-inflammatory cascade, in response to severe microbial infection or extensive tissue damage (such as caused by burns or multiple injuries)². Activation of the complement system and hyperactivation of cellular innate immune responses are associated with an excessive inflammatory response that characterizes sepsis. After being triggered by an overwhelming initial stimulus, neutrophils and macrophages produce and respond to cytokines, chemokines, complement-activation

products and other mediators. This pro-inflammatory environment causes the release of powerful secondary mediators (such as lipid factors and reactive oxygen species) that further amplify the inflammatory process. The malfunction of regulatory mechanisms during sepsis can result in a loss of control of inflammation, eventually leading to host damage due to overzealous activation of the inflammatory response.

However, the failure of anti-inflammatory therapies for sepsis in clinical trials raised the question of whether mortality in sepsis actually derives from an uncontrolled pro-inflammatory response³. Although some patients die during the initial, hyperinflammatory phase of sepsis, most patients succumb at later time points that are associated with a prolonged immunosuppressive state. Notably, neutrophils can undergo 'immune paralysis' during sepsis, which involves a complete shut-down of important intracellular signalling pathways, and dysfunction of the adaptive immune system is also an important contributing factor to the immunosuppression that is observed in the later stages of sepsis^{4,5}. T cells are thought to orchestrate the inflammatory response, particularly CD4⁺ T helper 1 (T_H1) cells and T_H2 cells, which have distinct cytokine profiles. During sepsis, the adaptive immune response diverts from an initial T_H1-cell response (characterized by interferon- γ (IFN γ) and interleukin-12 (IL-12) production) to a T_H2-cell response (characterized by IL-4, IL-5, IL-10 and IL-13 production), which can result in profound immunosuppression. The T_H1–T_H2-cell paradigm that describes how T_H cells interact with other immune cells has recently been expanded with the discovery of T_H17 cells, a subset of T_H cells that produces IL-17 (REF. 6). T_H17 cells are thought to be important for immunity to microorganisms that are not eliminated by T_H1- or T_H2-cell-mediated immune responses.

Department of Pathology,
The University of Michigan
Medical School, 1301
Catherine Road, Ann Arbor,
Michigan 48109-0602,
USA.

Correspondence to P.A.W.
e-mail: pward@umich.edu
doi:10.1038/nri2402
Published online
19 September 2008

Increased levels of apoptosis in lymphocytes and dendritic cells (DCs) further contribute to the suppression of immune responses during sepsis (reviewed in REF. 4). In addition to causing a marked decrease in cell numbers, the apoptosis of lymphocytes and DCs contributes to immunoparalysis through the immunosuppressive effects of apoptotic cells. However, different types of immune cell receive different apoptotic signals during sepsis. In contrast to lymphocytes and DCs, the apoptosis of macrophages and neutrophils seems to be unaffected or even decreased during sepsis^{7,8}. Whereas the increased apoptosis of lymphocytes and DCs results in severe immunosuppression, which places the patient at risk of nosocomial infections, decreased neutrophil apoptosis increases the bystander damage caused by their pro-inflammatory activity. Recent data indicate that T-cell-mediated suppression of the early innate immune response is required to minimize damage to the host and maximize the host defence response⁹.

There is now evidence that sepsis is a condition that affects not only the immune system but also other biological systems, such as the coagulation system and the autonomic nervous system (ANS)^{10–12}. In this Review, we describe the interplay between normally host-protective mechanisms that, through amplification or suppression during sepsis, can become instruments of harm. We discuss the mechanisms that initiate dysregulation of the inflammatory response and describe the role of specific inflammatory mediators that act as ‘central hubs’ to connect the various components of this response. In addition, we describe the pathogenic roles of the plasmatic cascades (the coagulation, fibrinolysis and complement systems) and the recently recognized interactions that occur between them, as well as new insights regarding the influence of the ANS on the inflammatory response. To illustrate the complexity of the inflammatory response in sepsis, we highlight the multidirectional interactions between the various systems that contribute to sepsis pathogenesis in a complex ‘inflammatory network’.

Initiation of the inflammatory response

Immune cells express a set of receptors known as pattern-recognition receptors (PRRs) that rapidly initiate host defence responses after detection of tissue damage or microbial infection. The presence of a microbial infection is detected by recognizing conserved pathogen-associated molecular patterns (PAMPs) that are expressed by both invading and innocuous microorganisms. By contrast, immune recognition of damaged tissue is mediated by intracellular proteins or mediators that are released from dying cells. These proteins are known as ‘alarmins’ and, together with PAMPs, are referred to as damage-associated molecular patterns (DAMPs)¹³. Toll-like receptors (TLRs), which are a subfamily of PRRs, have emerged as crucial receptors for the recognition of DAMPs and initiation of the inflammatory response.

During sepsis, there is a full-blown, systemic activation of immune responses due to the release of very high levels of DAMPs from invading microorganisms and/or damaged host tissue, which leads to the overstimulation

of immune cells. As a result, sepsis is accompanied by a markedly imbalanced cytokine response (known as a ‘cytokine storm’), which converts responses that are normally beneficial for fighting infections into excessive, damaging inflammation.

TLR4-mediated recognition of lipopolysaccharide (LPS), a well-characterized PAMP that is found in the outer membrane of Gram-negative bacteria, is thought to be an important trigger of the inflammatory response in sepsis^{14,15}. TLR4 forms a receptor complex with CD14 and MD2, the latter of which also has an important role in the recognition of LPS^{16,17}. In addition to LPS, various endogenous ligands for TLR4 have been described, including high-mobility group box 1 protein (HMGB1), which is an important mediator during the late phase of sepsis (see later)¹⁸. In the past, however, studies of TLR4 have been problematic owing to LPS contamination in recombinant proteins, which might limit the informative value of some studies.

It has been postulated that crosstalk occurs between TLR4 and the complement system, both of which are involved in the initiation of the inflammatory response in sepsis^{19,20}. Most strikingly, the complement anaphylatoxin C5a negatively regulates TLR4-mediated responses¹⁹. The extent of the regulatory effect of complement on TLR4-mediated cytokine production correlates with the level of complement-activation products and, in turn, the cytokines that are induced by TLR4 activation upregulate expression of the complement anaphylatoxin receptors C5AR and C3AR^{20,21}. The finding that activation of TLR4 in platelets initiates the formation of neutrophil extracellular traps to ensnare bacteria in the vasculature further demonstrates the intricate interplay between innate immunity and the clotting system in sepsis²².

Owing to its prominent role in the initiation of the inflammatory response, TLR4 is a potential therapeutic target for sepsis. In a recent study, antibody-mediated blockade of TLR4 and MD2 protected against polymicrobial sepsis²³ (TABLE 1). However, the mortality of septic mice that expressed a dysfunctional mutant TLR4 protein was not significantly different from that of wild-type mice with sepsis²⁴. In human sepsis, clinical trials that blocked TLR4 did not show beneficial effects²⁵, possibly because targeting TLR4 might only be an effective treatment for sepsis caused by Gram-negative bacteria or LPS, whereas the incidence of Gram-positive and fungal sepsis is increasing.

So, the early phase of sepsis, which is caused by the excessive activation of the host pathogen-recognition system by large-scale tissue damage and/or severe infection, leads to severe dysregulation of various body systems as a result of the pro-inflammatory environment.

Harmful central hubs in sepsis

The discovery that inflammatory mediators — not only invading microorganisms — are involved in the pathogenesis of sepsis has opened up a new avenue for the investigation of pathological mechanisms of inflammation. Many different mediators have been linked to the pathology of sepsis, some of which can be considered to be central hubs in the inflammatory

Nosocomial infections

These are infections that occur during treatment in a hospital or a healthcare service unit and are secondary to the patient's original condition. Nosocomial stems from the Greek word ‘nosokomeion’ meaning hospital (‘nosos’, disease; ‘komeo’, to take care of). This type of infection is also known as a hospital-acquired infection.

Anaphylatoxin

The pro-inflammatory complement-activation fragments C3a, C4a and C5a are also known as anaphylatoxins. They mediate inflammatory responses through cell activation and induce effects such as chemotaxis and histamine release.

Neutrophil extracellular traps

(NETs). A set of extracellular fibres produced by activated neutrophils to ensnare invading microorganisms. NETs enhance neutrophil killing of extracellular pathogens while minimizing damage to host cells.

Table 1 | Potential therapeutic targets in sepsis

System	Proposed mechanism	Target	References
Pattern-recognition system	Inhibition of PRRs to dampen the inflammatory response	TLR4 blockade	23
		RAGE blockade	49
Pro-inflammatory mediators	Blockade of central hubs of the inflammatory response to reverse established sepsis	MIF blockade or inhibition of its tautomerase activity	36,37
		HMGB1 blockade	44,50
		IL-17A blockade	52
Complement system	Neutralization of the harmful effects of C5a; formation of the MAC not affected	C5a neutralization	58
		Dual C5AR and C5L2 blockade	29
Coagulation system	Induction of anticoagulant and anti-inflammatory effects	Administration of activated protein C	74
		Selective PAR1 and PAR2 activation	83
Autonomic nervous system	Activation of the cholinergic anti-inflammatory pathway and/or suppression of the adrenergic pro-inflammatory pathway to restore homeostasis	Parasympathetic branch: • Pharmacological stimulation of α_7 nAChR on immune cells • Vagus-nerve stimulation	42 104
		Sympathetic branch: • Pharmacological modulation of α - and β -adrenergic receptor pathways on leukocytes	12, 108

α_7 nAChR, α_7 -nicotinic acetylcholine receptor; C5AR, C5a receptor; C5L2, C5a-like receptor 2; HMGB1, high-mobility group box 1 protein; IL-17A, interleukin-17A; MAC, membrane-attack complex; MIF, macrophage migration-inhibitory factor; PAR1, protease-activated receptor 1; PRR, pattern-recognition receptor; RAGE, receptor for advanced glycation end-products; TLR4, Toll-like receptor 4.

network (FIG. 1). Although they differ in terms of their source, kinetics of release and the stage of sepsis during which they predominate, these central hubs all have pleiotropic effects and connect various pathways of the immune response.

C5a. As part of the innate immune response, the complement system is activated during the early stages of sepsis, which generates large amounts of the anaphylatoxin C5a. At high concentrations, C5a has numerous harmful effects (reviewed in REF. 26; see later). Accordingly, C5a acts as a central mediator in sepsis by modulating other systems — including the coagulation cascade, TLR4-mediated responses and the release of cytokines, such as macrophage migration-inhibitory factor (MIF) and HMGB1 (REFS 10,19,27–29).

MIF. MIF, which was one of the first cytokines to be discovered, has a pivotal role in regulating systemic and local inflammatory responses (reviewed in REF. 30). Bacterial endo- and exotoxins, and pro-inflammatory mediators — such as tumour-necrosis factor (TNF), IFN γ and C5a — are strong inducers of MIF secretion by leukocytes^{28,30}. Unlike other cytokines, MIF is constitutively expressed by leukocytes and stored intracellularly³⁰. After its secretion, MIF functions as a classical pro-inflammatory cytokine and promotes innate and adaptive immune responses by activating macrophages and T cells³⁰. Interestingly, the pro-inflammatory activities of MIF are mediated by its tautomerase activity, which is encoded by a domain containing an evolutionarily conserved catalytic site³¹. In addition to mediating its own pro-inflammatory effects, MIF also induces and amplifies the production of other pro-inflammatory cytokines and upregulates the expression of TLR4 by phagocytes³⁰. At high concentrations, MIF prevents the

p53-dependent apoptosis of activated macrophages, which results in sustained inflammatory responses³⁰. However, the exact mechanisms by which MIF exerts its biological effects in the context of inflammation are not entirely clear. Although MIF activates intracellular signalling pathways after its endocytosis (which is an atypical mode of cytokine action)³⁰, the CD74 receptor complex has also been described to function as a MIF receptor, from which signals are transduced through CD44 (REF. 32).

MIF is unique among cytokines in that it links the immune system with the endocrine system. In response to stress, MIF is secreted by the hypothalamus, the anterior pituitary gland and the adrenal glands^{30,33}. Importantly, MIF antagonizes and overrides the anti-inflammatory effects of endogenous steroids³⁰, which might have consequences for the well-established use of corticosteroids as a therapy for sepsis. Endogenous corticosteroids induce the release of MIF from immune cells, and so the inhibitory effect of MIF on the action of corticosteroids is a negative-feedback loop^{30,34,35}. Excessive production of MIF is harmful in the acute phase of sepsis and plasma levels of MIF correlate with sepsis severity³⁶. Neutralization of MIF or targeting of its tautomerase activity attenuated the inflammatory response and improved survival in experimental sepsis^{36,37}. In addition, this treatment approach also markedly improved survival even when started after the onset of disease, which indicates that MIF is a promising therapeutic target^{36,37} (TABLE 1).

HMGB1. HMGB1 was originally described as a transcription factor³⁸. After its redefinition as a pro-inflammatory cytokine³⁹, HMGB1 became the focus of a large number of studies. HMGB1 is expressed by almost all cell types, except those lacking a nucleus

Tautomerase activity

The ability to catalyse the tautomerization (switching from one isomeric form to another) of D-dopachrome and L-dopachrome methyl ester into their corresponding indole derivatives. This reaction was used by early life forms for synthetic pathways. Macrophage migration-inhibitory factor (MIF) has been shown to have tautomerase activity; this evolutionarily conserved catalytic function is responsible for its pro-inflammatory effects.




Figure 1 | Central hubs of the inflammatory response in sepsis. During sepsis, the complement anaphylatoxin C5a is generated following the activation of the complement system and by the C5-convertase activity of thrombin of the coagulation cascade. C5a triggers the release of pro-inflammatory mediators, including macrophage migration-inhibitory factor (MIF) and high-mobility group box 1 protein (HMGB1), and it activates the coagulation cascade by inducing tissue-factor expression (not shown). HMGB1 is a pleiotropic cytokine that binds to Toll-like receptor 4 (TLR4) and acts as an endogenous alarmin to increase the release of pro-inflammatory mediators. TLR4-mediated responses, in turn, are negatively regulated by C5a. Similar to HMGB1, large amounts of MIF are released during sepsis, which promotes a pro-inflammatory response by amplifying cytokine secretion through the upregulation of TLR4 expression. MIF, which is produced by the pituitary gland as well as by leukocytes, inhibits the anti-inflammatory effects of endogenous glucocorticoids of the endocrine system, which, in turn, induce MIF secretion. HMGB1 links the immune response with the autonomic nervous system, which regulates the release of HMGB1 and other cytokines during sepsis. Interleukin-17A (IL-17A), which is an important regulator of inflammation at the interface between innate and adaptive immunity, orchestrates responses of both innate and adaptive immune cells.

Cholinergic anti-inflammatory pathway

This pathway fine-tunes cytokine production during inflammation in a highly regulated and reflexive manner. Interaction of acetylcholine with the α_7 -nicotinic acetylcholine receptor (α_7 nAChR) expressed by macrophages results in the suppression of pro-inflammatory cytokine production. The main component of this pathway is the vagus nerve of the parasympathetic branch of the autonomic nervous system.

(such as erythrocytes), and the main sources of HMGB1 in inflammation are macrophages, monocytes and neutrophils^{40,41}. HMGB1 can be secreted by immune cells after its acetylation in the nucleus and subsequent translocation to the cytoplasm, or it can be released from necrotic cells⁴⁰. The active secretion of HMGB1 is regulated by nuclear factor- κ B (NF- κ B) activation, probably through non-transcriptional mechanisms, although how this occurs is not well understood⁴². Intriguingly, although apoptotic cells are not a source of extracellular HMGB1 (REF. 43), they cause macrophages to release HMGB1 during sepsis⁴⁴. Extracellular HMGB1 specifically interacts with PRRs, including the receptor for advanced glycation end-products (RAGE), TLR2 and TLR4. HMGB1-induced signalling has pleiotropic effects on immune cells, promoting inflammation and the potentially harmful disruption of epithelial barriers^{18,40,45}. In addition to the activation of PRRs, HMGB1 increases the pro-inflammatory activity of cytokines (such as IL-1 β) through binding to these mediators, which supports the idea that HMGB1 might not act solely as a pro-inflammatory mediator, but might also function as a carrier or DAMP^{46,47}.

Although HMGB1 is released systemically during sepsis, plasma levels do not necessarily correlate with outcome or survival⁴⁰. In contrast to other sepsis-associated cytokines, the peak of HMGB1 release occurs during later stages of the disease, and the levels of HMGB1 do not always decrease in patients who have recovered from sepsis^{39,48}. Pathogen-derived molecules and pro-inflammatory stimuli (such as TNF, IL-1 β and IFN γ) induce the secretion of HMGB1 during inflammation⁴⁰. Interaction between C5a and its other receptor, C5a-like receptor 2 (C5L2), also triggers the release of HMGB1 in sepsis²⁹. Interestingly, HMGB1 secretion is under the influence of the ANS⁴². Activation of the cholinergic anti-inflammatory pathway suppresses HMGB1 secretion by macrophages in sepsis and improves survival⁴².

Owing to the pleiotropic effects of HMGB1 on the inflammatory response and its late release in sepsis, targeting HMGB1 might be a promising therapeutic strategy. In experimental settings, the direct blockade of HMGB1 or inhibition of RAGE improved survival in endotoxaemia and experimental sepsis^{39,44,49,50} (TABLE 1). Similar to the targeting of MIF, neutralization of HMGB1 prevented lethality when treatment occurred after the onset of sepsis and it reversed the development of multi-organ failure^{44,50} (TABLE 1). However, the complexity of the underlying mechanisms of HMGB1 function precludes the use of HMGB1 blockade in clinical trials at this point.

IL-17A. The recent discovery of the IL-17 cytokine family, the members of which have emerged as important mediators of immune regulation, has greatly improved our understanding of the interplay between innate and adaptive immune responses (reviewed in REF. 51). IL-17A, the first described member of the IL-17 family, is a pro-inflammatory cytokine that is mainly produced by T_H17 cells⁵¹. IL-17A is also secreted by various other types of immune cell, including neutrophils, CD8⁺ T cells, natural killer cells, other T_H-cell subsets and $\gamma\delta$ T cells⁵¹. In brief, IL-17A is involved in mediating pro-inflammatory responses by triggering the production of many other cytokines (such as IL-1 β , IL-6 and TNF) and provides crosstalk between lymphocytes and phagocytes⁵¹.

It has recently been shown that increased IL-17A levels have adverse effects during experimental sepsis⁵². Neutralization of IL-17A markedly improved survival, even when the treatment was administered as late as 12 hours after the initiation of experimental sepsis⁵² (TABLE 1). The protective effects of IL-17A blockade were associated with a marked attenuation of bacteraemia and markedly decreased plasma levels of pro-inflammatory cytokines⁵². In agreement with these data, the *in vitro* production of pro-inflammatory mediators by macrophages in response to LPS was significantly increased in the presence of recombinant IL-17A⁵². However, it is not yet known whether levels of IL-17A are increased in patients with sepsis, or during which phase of sepsis the neutralization of IL-17A would be beneficial in the clinical setting. Because the production of IL-17 is important for directing the immune response against some specific infections, the

Endotoxaemia

This is caused by the presence of endotoxins, which are derived from Gram-negative bacteria, in the blood. It results in systemic activation of the inflammatory response, the development of shock and multi-organ failure and death. Models of endotoxaemia are used in experimental settings to induce systemic inflammation, but they do not necessarily mimic human sepsis.

Septic cardiomyopathy

The decreased myocardial function that occurs during sepsis-associated multi-organ failure. Hypotheses concerning the aetiology of this decreased function include impairment of mitochondrial function, dysfunction of the β -adrenoceptor-G-protein-adenylate cyclase system, calcium-channel blockade by direct and indirect cardiodepressant factors and contractile impairment by activated leukocytes.

blockade of IL-17A under certain conditions might cause more harm than good. Therefore, it remains to be determined whether IL-17A is a useful target for therapeutic intervention in sepsis.

Dysregulation of plasmatic cascades

Complementopathy. The complement system can be activated through three different pathways, which converge on the generation of the anaphylatoxins C3a and C5a, C4a and the membrane-attack complex (MAC; also known as C5b-C9). In clinical studies of sepsis, increased concentrations of C3a, C4a and C5a in the plasma have been linked to poor outcome and survival^{53,54}. Interestingly, C3a might have anti-inflammatory properties in addition to acting as a pro-inflammatory anaphylatoxin⁵⁵. Mice with C3AR deficiency were more susceptible to endotoxin shock, which was accompanied by an increase in the concentration of pro-inflammatory cytokines in the plasma. Binding of C3a to C3AR can trigger the secretion of anti-inflammatory hormones by the pituitary gland⁵⁶, which might account for the anti-inflammatory properties of C3a.

New discoveries continue to increase our understanding of the numerous harmful effects of excessive C5a production during sepsis. The effects of C5a contribute to immunoparalysis⁵⁷, multi-organ failure⁵⁸, the apoptosis of thymocytes⁵⁹ and adrenal medullary cells⁶⁰, and imbalances in the coagulation system⁶¹ (FIG. 2). In addition, C5a

is involved in the development of septic cardiomyopathy⁶². Decreased pressure in the left ventricle of the heart has been observed following sepsis, accompanied by defective contractility of cardiomyocytes; both of these effects were reversed by administration of a C5a-specific blocking antibody⁶². Furthermore, when recombinant C5a was added to isolated rat cardiomyocytes *in vitro*, contractile dysfunction was induced⁶², which indicates that excessive generation of C5a during sepsis causes cardiomyopathy.

Recent studies corroborate an important role for C5a in sepsis and indicate that it exerts its harmful effects in a complex manner²⁹. In addition to C5AR, C5a can bind specifically to a second receptor, C5L2, the function of which was unknown until recently. It was originally hypothesized that C5L2 functions as a decoy receptor for C5a, competing with C5AR for binding of C5a⁶³, although newer evidence indicates that C5L2 is a functional receptor^{29,64}. In human sepsis, the expression of C5L2 was downregulated on the surface of neutrophils during septic shock⁶⁵. The extent of this downregulation correlated with the development of multi-organ failure, which indicates that C5L2 contributes to the pathogenesis of sepsis⁶⁵. There is now evidence that C5AR and C5L2 cooperatively enhance the inflammatory response during sepsis, although each receptor might have specific and distinct functional roles²⁹. For example, C5a-induced release of MIF depends on C5AR signalling, whereas C5L2 mediates the C5a-dependent release of




Figure 2 | C5a is a central mediator of the inflammatory response in sepsis. During the early stages of sepsis, the complement system is systemically activated, generating large amounts of the anaphylatoxin C5a. C5a, which is a central molecule in the immunopathogenesis of sepsis, exerts its effects through interactions with its two receptors, C5a receptor (C5AR) and C5a-like receptor 2 (C5L2). The expression of these receptors is upregulated during sepsis, and their interactions with C5a contribute synergistically to harmful events in sepsis. The numerous effects of C5a include activation of the coagulation cascade by the induction of tissue-factor expression, which can result in disseminated intravascular coagulation (DIC). Furthermore, C5a triggers the release of pro-inflammatory cytokines, including macrophage migration-inhibitory factor (MIF) and high-mobility group box 1 protein (HMGB1), which contribute to the systemic inflammatory response syndrome (SIRS). In the later stages of sepsis, C5a is also responsible for sepsis-induced neutrophil dysfunction, leading to the shut down of intracellular signalling (immune paralysis) and increased susceptibility to secondary infections. C5a-induced apoptosis of thymocytes further aggravates immunosuppression, whereas the apoptosis of adrenal medullary cells results in insufficiency of the adrenergic system, eventually leading to septic shock. Recently, C5a and C5AR were also shown to be directly involved in the development of septic cardiomyopathy.

Disseminated intravascular coagulation

(DIC). Also known as consumptive coagulopathy, this is a pathological process in which the blood begins to coagulate throughout the entire body. During this process, platelets and coagulation factors are depleted, resulting in a paradoxical situation in which there is a high risk of simultaneous fatal thrombosis and large-scale haemorrhage. DIC often occurs in critically ill patients with overwhelming infection, fulminant sepsis or malignancy.

Thrombin

Thrombin (also known as activated Factor II) is the central serine protease that converts soluble fibrinogen into insoluble strands of fibrin. It also catalyses many other coagulation-related reactions.

Tissue factor

A pro-coagulant factor that stimulates thrombus formation following contact with blood by accelerating the action of the coagulation factors Factor VIIa and Factor Xa. It can also be expressed on the surface of activated endothelial cells.

Activated protein C

A physiological anticoagulant. The activated form degrades Factor Va and Factor VIIIa of the coagulation cascade. The protein-C pathway has anti-thrombotic activity, as well as anti-inflammatory and anti-apoptotic functions. Administration of human recombinant activated protein C for the treatment of sepsis might block dysregulated coagulation, inhibit pro-inflammatory pathways and preserve organ function.

Thrombomodulin

An integral membrane protein that is expressed on the surface of endothelial cells. It functions as a co-factor in thrombin-induced activation of protein C in the anticoagulant pathway by forming complexes with thrombin. Thrombomodulin–thrombin complexes also stimulate fibrinolysis by cleaving thrombin-activatable fibrinolysis inhibitor (TAFI) into its active form.

HMGB1 from phagocytes^{28,29}. Importantly, blockade of either receptor protects against lethality in moderate forms of sepsis, but only the inhibition of both C5a receptors provides protection in severe sepsis²⁹.

Although inhibitors of other complement factors (such as a C1 inhibitor or soluble recombinant complement receptor 1) have had limited beneficial effects in the clinic^{66,67}, C5a might be a promising target for pharmaceutical intervention in sepsis. The advantage of this strategy is that the inhibition of the harmful effects of C5a does not interfere with the assembly of the MAC, which is essential for defence against invading microorganisms⁶⁸. Currently, dual blockade of C5AR and C5L2, rather than blockade of C5a alone, seems to be an encouraging strategy for clinical trials²⁹ (TABLE 1). However, as complement activation is an early event in sepsis, the availability of a reliable and sensitive bedside test to assess the extent of complement activation in a patient will be essential for successful intervention directed at the complement cascade.

Coagulopathy. In the clinical setting of sepsis, dysregulation of the coagulation cascade (BOX 1) results in major complications⁶⁹. The extent of activation of the coagulation cascade during sepsis can range from an insignificant level to the occurrence of disseminated intravascular coagulation (DIC). In the initial phase of DIC, thrombin activation results in intra- and extravascular fibrin formation (a process known as hypercoagulability), followed by the consumption of coagulation factors and platelet dysfunction (known as hypocoagulability)⁷⁰. In the late phase of DIC, microvascular fibrin deposition is often associated with the development of multi-organ failure owing to perturbations in the microcirculation⁷⁰. As DIC develops, inflammation and coagulation interact in a bidirectional manner⁷¹. Activated **thrombin** can promote the activation of various pro-inflammatory pathways — including the

production of pro-inflammatory cytokines (such as TNF, IL-1 β and IL-6) and the generation of C5a — and cytokines, in turn, can stimulate coagulation^{10,72–74}. Tissue factor, which is a central molecule in the initiation of DIC, is expressed by activated endothelial cells and by cells that are not normally exposed to blood flow, such as sub-endothelial cells and fibroblasts, and also by circulating immune cells⁷⁵. In sepsis, the pro-inflammatory environment causes mononuclear cells to upregulate the expression of **tissue factor** on their cell surface, leading to the systemic activation of coagulation⁷⁶.

Another consequence of DIC is the inhibition of fibrinolysis. In addition to endothelial-cell dysfunction during sepsis, which also occurs as a result of the pro-inflammatory environment, increased levels of plasminogen-activator inhibitor 1 (PAI1) and thrombin-activatable fibrinolysis inhibitor (TAFI) lead to impaired removal of fibrin⁷⁷. Also, the consumption of various factors that normally regulate the generation of thrombin, such as **antithrombin III**, **protein C** and **tissue-factor pathway inhibitor (TFPI)**, contributes to the development of DIC⁷⁸.

Recombinant activated protein C is currently the only approved drug for the treatment of sepsis that targets the inflammatory response⁷⁴ (TABLE 1). Protein C, which is a regulator of the coagulation cascade, is activated by thrombin bound to **thrombomodulin** and by endothelial protein C receptor (EPCR) on endothelial-cell membranes⁷⁹. After dissociation from EPCR, activated protein C binds to its co-factor, **protein S**, which then results in the inactivation of clotting factors Va and VIIIa⁷⁹. In addition to its anticoagulant activity, activated protein C has profound anti-apoptotic and anti-inflammatory properties. It markedly decreases the apoptosis of endothelial cells and lymphocytes and exerts profibrinolytic effects by inhibiting PAI1 (REFS 74,80). The anti-inflammatory effects of activated protein C are

Box 1 | Coagulation

The coagulation cascade is initiated by the exposure of coagulation factors in the blood to subendothelial proteins following damage to the blood-vessel endothelium. In primary haemostasis, circulating platelets bind to collagen through their cell-surface glycoprotein Ia/IIa receptors to form a haemostatic plug at the site of injury. The adhesion of platelets is stabilized by large, multimeric von-Willebrand-factor proteins, which form links between platelets, glycoproteins and collagen fibrils. Simultaneously, the action of a complex cascade of coagulation factors (a group of serine proteases that are activated in a sequential manner) results in the formation of fibrin strands, which further strengthen the platelet plug (secondary haemostasis). Traditionally, the coagulation cascade has been described as two pathways: the contact-dependent (intrinsic) activation pathway and the tissue-factor (extrinsic) pathway, the latter being the main pathway for the initiation of blood coagulation. These two pathways converge on the activation of thrombin, which converts fibrinogen to fibrin and ultimately results in the formation of a fibrin-crosslinked clot. The contemporary description of physiological haemostasis *in vivo* does not divide coagulation into cellular and plasmatic components or different activation pathways, but instead describes that coagulation involves three phases¹²¹. First, the initiation phase is characterized by the exposure of tissue factor after endothelial damage, resulting in the activation of thrombin. Second, thrombin augments coagulation by fully activating platelets and increasing platelet adhesion during the amplification phase. Third, large amounts of thrombin are generated on the surface of activated platelets, resulting in the stabilization of the blood clot in the propagation phase¹²¹.

Eventually, blood clots are organized (which involves the laying down of collagen and the formation of vascular channels) or absorbed by fibrin degradation (a process known as fibrinolysis). The main protease of the fibrinolysis cascade is plasmin, which is activated by tissue plasminogen activator, urokinase-like plasminogen activator, thrombin and fibrin itself. Under normal conditions, the balance between the coagulation and fibrinolysis systems, which is maintained by various regulatory mechanisms, prevents intravascular coagulation.

mediated by EPCR and the cleavage of protease-activated receptor 1 (PAR1), which has a central role in linking coagulation and inflammation⁸¹⁻⁸³ (TABLE 1).

The protein-C pathway is particularly susceptible to inhibition by inflammatory responses in sepsis-associated DIC⁷¹. In addition to a decrease in the level of protein C, the downregulation, shedding and cleavage of thrombomodulin and EPCR are the main causes of dysfunction of the protein-C pathway⁸⁴. HMGB1 inhibits the protein-C pathway by interfering with the thrombin-thrombomodulin complex and it also promotes coagulation by stimulating tissue-factor expression and inhibiting tissue plasminogen activator (TPA), a serine protease on the surface of endothelial cells that activates plasmin of the fibrinolysis cascade⁸⁵.

The administration of activated protein C in sepsis suppresses pro-inflammatory cytokine production and decreases the adhesion of phagocytes to injured endothelium through EPCR- and PAR1-dependent signalling^{74,86}. However, the anticoagulant activity of activated protein C might exacerbate bleeding complications in patients who usually have a compromised clotting system⁷⁴. Future clinical trials should assess whether a form of activated protein C that does not have anticoagulant effects⁸⁷ will improve clinical efficacy (that is, decrease sepsis mortality) and safety (that is, decrease the incidence of bleeding complications) in humans with sepsis.

Linking complement and coagulation. Traditionally, the complement and coagulation systems are described as separate cascades. As descendants of a common ancestral pathway, both are proteolytic cascades that are composed of serine proteases with common structural characteristics and similar activating stimuli^{71,88}. The relationship is not limited to the biochemical similarity in their serine proteases, however, as these two pathways are also linked by many mutual connections that form a complex network (FIG. 3).

During sepsis, the activated coagulation pathway predisposes to thrombosis and DIC, which further aggravate the excessive inflammatory response and complement activation⁷¹. A well-known interaction between the complement and coagulation systems is the activation of the classical complement pathway by coagulation Factor XIIa, which can activate the complement component C1 (REF. 89). More recently, it has been shown that thrombin can function as a C5 convertase in a C3-independent manner¹⁰. This crosstalk is particularly interesting, not only because thrombin and C5a are central factors in their respective cascades, but also because this indicates that C5a and the MAC can be generated in the absence of upstream complement activation. Similar to thrombin, kallikrein and plasmin directly cleave C3 and its activation fragments^{90,91}. In an indirect negative-feedback loop, thrombin-activated TAFI inactivates C3a and C5a⁹².




Figure 3 | Cross-talk between the complement, coagulation and fibrinolysis systems. The complement system, the coagulation cascade and the fibrinolysis cascade communicate through many direct and bidirectional interactions (indicated by red arrows). Activated clotting Factor XII (FXIIa) can activate the classical complement pathway through cleavage of the complement component C1. Similarly, thrombin, kallikrein (not shown) and plasmin directly cleave complement component C3, as well as its activation fragments. Moreover, thrombin can cleave C5 into C5a, which occurs independently of C3 and therefore represents a bypass of the three traditional complement-activation pathways (that is, the classical, lectin and alternative pathways). Thrombin-activatable fibrinolysis inhibitor (TAFI) inactivates C3a and C5a in a negative-feedback loop. The complement system also amplifies coagulation through the C5a-mediated induction of expression of tissue factor and plasminogen-activator inhibitor 1 (PAI1) by leukocytes, the latter of which inhibits fibrinolysis. In addition, mannan-binding lectin serine protease 2 (MASP2) of the lectin complement-activation pathway triggers coagulation by converting prothrombin to thrombin. C4b-binding protein (C4BP) of the complement pathway inhibits protein S, which is a co-factor for the activated protein-C pathway of coagulation inhibition, which indicates that the inhibition of anticoagulant mechanisms further augments the pro-coagulant activities of complement. MAC, membrane-attack complex (C5b-C9); TPA, tissue plasminogen activator; UPA, urokinase-like plasminogen activator.

Box 2 | The autonomic nervous system

The autonomic nervous system (ANS) is part of the peripheral nervous system. It has three components: the parasympathetic branch, the sympathetic branch and the enteric nervous system. The ANS maintains homeostasis in the body by controlling vital functions that include heart rate, respiration rate, digestion, perspiration and body temperature. Traditionally, the sympathetic and parasympathetic branches of the ANS were thought to be endogenous neuronal antagonists. Therefore, the classical terminology referred to adrenergic responses (sympathetic) as 'fight or flight' and to cholinergic responses (parasympathetic) as 'rest and digest' responses. However, it is now clear that the relationship between these pathways is more complex.

Signal transmission in the parasympathetic branch of the ANS is mediated by acetylcholine and its receptors, which are abundantly expressed by many cell types¹²². The main peripheral component of the parasympathetic branch of the ANS is the vagus nerve.

The sympathetic branch of the ANS consists of sympathetic neurons and the adrenal medulla. Catecholamines, which are the main mediators of the sympathetic branch, mediate pleiotropic effects by interacting with adrenergic receptors that are ubiquitously expressed by nearly all tissue and cell types¹²². The activation of adrenergic receptors triggers an intricate intracellular signalling network that has yet to be fully understood.

The enteric nervous system, which controls the gastrointestinal system, is also considered to be part of the ANS. Anatomically, the enteric nervous system consists of a large number of neurons that are embedded in the lining of the gastrointestinal tract. Although the enteric nervous system can operate autonomously, it communicates closely with the central nervous system, and is associated with a considerable amount of sympathetic and parasympathetic innervation.

The complement system amplifies coagulation by modification of phospholipid membranes (which is required for the initiation of coagulation through tissue factor), by activating platelets and by inducing the expression of tissue factor and PAI1 by leukocytes^{71,93,94}. Accordingly, blockade of C5a during experimental sepsis markedly ameliorated the effects of DIC⁶¹. In addition, mannan-binding lectin serine protease 2 (MASP2), a protease that is characteristic of the lectin pathway of complement activation, can activate coagulation by cleaving prothrombin into active thrombin⁹⁵. The pro-coagulant activities of complement are increased when anticoagulant mechanisms are inhibited; for example, the formation of a complex between C4b-binding protein and protein S results in a decrease in the availability of protein S to act as a co-factor for the anticoagulant protein-C pathway⁹⁶. In addition, several indirect influences of the complement system on coagulation that are mediated through other pro-inflammatory factors (such as TNF, IL-6 and HMGB1) have been documented.

In summary, the complement, coagulation and fibrinolysis systems are tightly connected through multiple direct interactions of serine proteases, which together make up a 'plasma serine-protease network'. In the setting of sepsis, crosstalk between the complement and coagulation pathways is of particular importance, as their uncontrolled activation is an essential contributor to the pathogenesis of the disease.

The autonomic nervous system

Recent advances in the field of neuroimmunology have shown that the nervous system and the immune system communicate during inflammation. The main pathways involved in this crosstalk are the hypothalamic–pituitary–adrenal axis and the ANS^{11,97,98} (BOX 2). Immune cells can also synthesize and release neurotransmitters and express receptors for these mediators^{11,12,99}. So, these neuromediators function as the biochemical language of the neuro–endocrine–immune network, which allows the body to adapt rapidly to changes of internal and external environments. Accordingly, Munford and Tracey have suggested that severe sepsis is a neuro–endocrine disorder¹⁰⁰.

The parasympathetic ANS. Signalling of the vagus nerve by engagement of cholinergic receptors expressed by phagocytes has an important regulatory role in inflammation¹⁰¹. Activation of α_7 -nicotinic acetylcholine receptors (α_7 nAChRs), either by vagus-nerve stimulation or by α_7 nAChR agonists, decreases intracellular cytokine synthesis by macrophages and dampens the inflammatory response^{42,101} (FIG. 4). In other words, inflammation is under neuronal control by the ANS, which can reflexively modulate the inflammatory response by inhibiting the production of pro-inflammatory cytokines. This concept has therefore been termed the 'inflammatory reflex'¹¹. The efferent arm of the inflammatory reflex is the cholinergic anti-inflammatory pathway¹⁰², which is a robust regulator of cytokine production.

Recent studies have shown that the branch of the vagus nerve that innervates the spleen is crucial for the suppression of cytokine synthesis in sepsis¹⁰³. The spleen is an important source of TNF during sepsis and splenectomy significantly decreases systemic and hepatic levels of TNF in septic mice^{102,103}. In addition, direct cholinergic modulation of immune cells that transit through the spleen contributes to the regulation of inflammation at distant sites. In experimental sepsis, activation of the cholinergic anti-inflammatory pathway inhibited the production of pro-inflammatory mediators (including HMGB1 and TNF) and markedly increased survival, even when carried out as late as 24 hours after the onset of disease^{42,104}.

The sympathetic ANS. The main transmitters of the sympathetic branch of the ANS are catecholamines, which act through binding to adrenergic receptors. The early phase of sepsis is characterized by high concentrations of circulating catecholamines, which boost the initial inflammatory response. Later in septic shock, the production and release of endogenous catecholamines can become insufficient for maintaining equilibrium of the cardiovascular system (as indicated by the need for catecholamine administration during septic shock). The depletion of endogenous catecholamine sources might be caused by the apoptosis of adrenal medullary cells⁶⁰.

Catecholamines

Tyrosine-derived mediators that are produced mainly by the adrenal medulla and by the postganglionic fibres of the sympathetic nervous system. Recently, it has been found that immune cells are also a source of catecholamines. The most abundant catecholamines are the biogenic amines adrenaline, noradrenaline and dopamine, which function as neurotransmitters in the sympathetic branch of the autonomic nervous system through interaction with adrenergic receptors expressed by numerous cell and tissue types.




Figure 4 | Effects of pathways of the ANS on inflammation during sepsis. The balance between the two branches of the autonomic nervous system (ANS) can direct the inflammatory response towards pro- or anti-inflammatory outcomes. Whereas activation of the cholinergic anti-inflammatory pathway (part of the parasympathetic branch of the ANS) dampens inflammation, stimulation of the adrenergic pathways leads to amplification of the inflammatory response. **a** | In the adrenergic pro-inflammatory pathway, high concentrations of circulating catecholamines amplify the initial inflammatory response, particularly in the early phase of sepsis. Sources for catecholamine production and release are the adrenal medulla, sympathetic neurons and leukocytes (phagocytic cells and lymphocytes). Catecholamines exert their immunomodulatory effects through α - and β -adrenergic receptors that are expressed by various cell types, resulting in the increased release of pro-inflammatory mediators. **b** | By contrast, the activation of the cholinergic anti-inflammatory pathway in sepsis attenuates the inflammatory response. These effects are mediated through engagement of α_7 -nicotinic acetylcholine receptors (α_7 nAChRs). Acetylcholine is released following vagus-nerve stimulation, resulting in inhibition of the synthesis and release of pro-inflammatory mediators such as high-mobility group box 1 protein and tumour-necrosis factor.

The concomitant dysfunctional adrenergic modulation of heart and blood vessels during septic shock indicates that impairment of adrenergic regulation contributes to cardio-circulatory failure¹⁰⁵.

It was originally thought that the synthesis of catecholamines was carried out only by the neuronal cells of the sympathetic branch of the ANS, but it has now been shown that leukocytes are also an abundant source of catecholamines^{12,99}. Leukocytes also express adrenergic receptors, which indicates that catecholamines might have autocrine and/or paracrine effects on immune cells^{12,99}. The activation of adrenergic receptors on immune cells triggers distinct and finely tuned cytokine responses through NF- κ B-dependent mechanisms^{12,99}.

During sepsis, catecholamines exert immunomodulatory effects through α - and β -adrenergic receptors that are expressed by immune cells^{106–108}. Stimulation of these receptors alters lymphocyte trafficking, vascular perfusion and cell proliferation and apoptosis, thereby affecting the functional responses of leukocytes^{109–111}. The response of neutrophils and macrophages in particular underlies adrenergic regulation by catecholamines, as the release of pro-inflammatory cytokines

by these cells is tightly regulated by α -adrenergic receptors^{12,111}. Catecholamines might also contribute to the deleterious effects of sepsis through direct stimulation of bacterial growth in the gastrointestinal system, which might contribute to bacteraemia through the translocation of enteric bacteria into the lymphatic and blood compartments¹¹².

In summary, activation of the adrenergic pathways of the sympathetic branch of the ANS during the early phase of sepsis promotes pro-inflammatory responses and aggravates adverse events, although the mechanisms that underlie these effects have yet to be evaluated in detail.

The enteric nervous system. Sepsis can occur as a result of inflammation in the abdominal cavity (known as peritonitis) that results from the disruption of barriers that protect the sterile compartment of the abdominal cavity from pathogens in the intestinal lumen. It was hypothesized that the translocation of enteric bacteria into the blood might also occur when the gut was not the main source of inflammation (for example, in situations such as pneumonia or burn injury), due to a general loss of intestinal epithelial barrier functions as a result of the pro-inflammatory environment. Recent research has shown that the gut can produce large amounts of catecholamines during sepsis, which are released into intestinal blood¹¹³. Catecholamines that drain from the intestines through the portal vein into the liver can alter the functional state of Kupffer cells and hepatocytes through α_2 -adrenergic receptor signalling¹¹⁴, ultimately contributing to the release of pro-inflammatory cytokines, hepatocellular dysfunction and liver failure^{113,115}. Catecholamine-induced activation of Kupffer cells might also be an important source of the cytokine storm during sepsis¹¹⁶. Additional studies are required to identify the cell source of intestine-derived catecholamines during sepsis. It remains possible that either the enteric nervous system or resident immune cells in Peyer's patches and lymph nodes of the intestinal system (or both) are responsible for the generation of intestine-derived catecholamines during sepsis.

ANS-targeted therapy. Modulation of the ANS might be a promising approach for the treatment of sepsis as an alternative to blocking pro-inflammatory mediators directly (TABLE 1). Whereas the cholinergic pathway attenuates the immune response and is considered to be anti-inflammatory¹¹, adrenergic stimulation promotes the release of pro-inflammatory mediators and the recruitment of leukocytes¹² (FIG. 4). Given that an imbalance between these two branches of the ANS contributes to the development of sepsis, stimulation of the cholinergic vagus nerve and/or suppression of adrenergic pathways might help to restore homeostasis. It has recently been shown that transcutaneous electrical stimulation of the vagus nerve improved survival and decreased the level of pro-inflammatory mediators in experimental sepsis¹⁰⁴. Although this approach might not yet be ready for use in the clinic, the fact that this treatment for sepsis would be non-invasive and independent of pharmacokinetics — unlike the administration of drugs — makes it particularly attractive.

Kupffer cells

The resident macrophages of the liver, which are derived from blood monocytes. They phagocytose pathogenic particles and microorganisms that have entered the liver sinusoids.




Figure 5 | The inflammatory network in sepsis. During sepsis, homeostasis between the various biological systems of the inflammatory network is highly imbalanced. In the initiation of sepsis, the release of a large amount of damage-associated molecular patterns (DAMPs) from invading microorganisms and/or damaged host tissue results in the overstimulation of pattern-recognition receptors (PRRs) on immune cells. Activated immune cells release excessive amounts of pro-inflammatory mediators (resulting in a 'cytokine storm'), free radicals and enzymes, which converts the normally beneficial effects of inflammation into an excessive response that damages the host. Activation of the adrenergic branch of the autonomic nervous system (ANS) and/or decreased activity of the cholinergic anti-inflammatory pathway (of the parasympathetic branch of the ANS) further amplifies the pro-inflammatory responses of neutrophils, macrophages and dendritic cells in sepsis. The presence of invading microorganisms or their products in the blood can cause systemic activation of the complement system, which results in the excessive generation of complement anaphylatoxins, which, at high concentrations, induce numerous harmful effects. Simultaneous activation of the coagulation system and the inhibition of fibrinolysis as a result of the pro-inflammatory environment and/or damaged endothelium can result in disseminated intravascular coagulation (DIC), which is a major complication of sepsis, and in the amplification of the inflammatory response. The complement, coagulation and fibrinolysis systems are tightly connected through direct interactions of serine proteases, and imbalances in each cascade are intensified in a positive-feedback loop (FIG. 4). Finally, the sustained pro-inflammatory environment affects the functional state of immune effector cells, eventually causing the dysfunction of neutrophils and immunoparalysis. Alterations in leukocyte apoptosis in the later stages of sepsis further account for immunosuppression, which increases the susceptibility to secondary infections.

Concluding remarks

Despite more than 20 years of extensive research, none of the promising therapeutic approaches for sepsis that target the inflammatory response has been successfully translated to the clinical setting, and rates of sepsis mortality have not decreased¹¹⁷. These failures are in part due to the fact that some of the animal models for sepsis that are used, such as endotoxaemia, do not accurately mimic sepsis in humans and/or because the limitations of animal models have been disregarded (reviewed in REF. 118). In addition, it is now clear that the belief that a single key mediator causes sepsis, and that neutralization of such a factor could be a cure for all patients with sepsis, is erroneous. Instead, we now understand that sepsis is a complex, dynamic syndrome with great heterogeneity, and not a distinct disease. Sepsis can result from various causative insults, and susceptibility can be influenced by premorbid factors that include ethnicity, gender, age, genetic defects and environmental factors¹¹⁹. In particular, genetic and epigenetic changes, such as mutations in genes that encode PRRs or mediators of inflammation and their receptors, might have consequences for the host response. Because the underlying inflammatory response during sepsis varies between individual patients, various options for therapy should be made available. Ideally, individual patients should be precisely monitored for changes in characteristic markers of the host immune response to aid in the choice of specific immunomodulatory therapies.

We now know that the underlying inflammatory response in sepsis involves a complex interplay of different biological systems and cell types, resulting in severe dysregulation of the inflammatory network (FIG. 5). We are just beginning to understand the underlying regulatory pathways of this network. The application of interdisciplinary approaches will improve our knowledge of the molecular biology of inflammation in the context of sepsis. Initial large-scale programs that aim to uncover the molecular mechanisms of inflammation — which include the fields of surgery, genomics, proteomics, biostatistics, bioinformatics, computational biology and genetics — are now underway¹²⁰. Although our current knowledge regarding inflammation might be in its infancy, recent progress indicates that patients will one day benefit from more advanced knowledge of the inflammatory response in sepsis.

1. Thomas, L. Germs. *N. Engl. J. Med.* **287**, 553–555 (1972).
2. Bone, R. C. *et al.* Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest* **101**, 1644–1655 (1992).
3. Remick, D. G. Cytokine therapeutics for the treatment of sepsis: why has nothing worked? *Curr. Pharm. Des.* **9**, 75–82 (2003).
4. Hotchkiss, R. S. & Nicholson, D. W. Apoptosis and caspases regulate death and inflammation in sepsis. *Nature Rev. Immunol.* **6**, 813–822 (2006).
5. Solomkin, J. S., Jenkins, M. K., Nelson, R. D., Chenoweth, D. & Simmons, R. L. Neutrophil dysfunction in sepsis. II. Evidence for the role of complement activation products in cellular deactivation. *Surgery* **90**, 319–327 (1981).
6. Harrington, L. E. *et al.* Interleukin-17-producing CD4⁺ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nature Immunol.* **6**, 1123–1132 (2005).
7. Hotchkiss, R. S. *et al.* Accelerated lymphocyte death in sepsis occurs by both the death receptor and mitochondrial pathways. *J. Immunol.* **174**, 5110–5118 (2005).
8. Hotchkiss, R. S. *et al.* Depletion of dendritic cells, but not macrophages, in patients with sepsis. *J. Immunol.* **168**, 2493–2500 (2002).
9. Kim, K. D. *et al.* Adaptive immune cells temper initial innate responses. *Nature Med.* **13**, 1248–1252 (2007).
10. Huber-Lang, M. *et al.* Generation of C5a in the absence of C3: a new complement activation pathway. *Nature Med.* **12**, 682–687 (2006). **This study shows that thrombin can act as a C5 convertase, which indicates that there is direct crosstalk between the coagulation and complement systems.**
11. Tracey, K. J. The inflammatory reflex. *Nature* **420**, 853–859 (2002).
12. Flierl, M. A. *et al.* Phagocyte-derived catecholamines enhance acute inflammatory injury. *Nature* **449**, 721–725 (2007). **This publication identifies phagocytes as a source of catecholamines, which enhance the inflammatory response.**
13. Bianchi, M. E. DAMPs, PAMPs and alarmins: all we need to know about danger. *J. Leukocyte Biol.* **81**, 1–5 (2007).
14. Medzhitov, R. & Janeway, C. Jr. Innate immunity. *N. Engl. J. Med.* **343**, 338–344 (2000).
15. Poltorak, A. *et al.* Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: mutations in *Tlr4* gene. *Science* **282**, 2085–2088 (1998). **In this study, TLR4 is defined as the recognition receptor for LPS.**
16. Ohto, U., Fukase, K., Miyake, K. & Satow, Y. Crystal structures of human MD-2 and its complex with antiendotoxic lipid IVa. *Science* **316**, 1632–1634 (2007).

17. Kim, H. M. *et al.* Crystal structure of the TLR4–MD-2 complex with bound endotoxin antagonist Eritoran. *Cell* **130**, 906–917 (2007).

18. Park, J. S. *et al.* Involvement of Toll-like receptors 2 and 4 in cellular activation by high mobility group box 1 protein. *J. Biol. Chem.* **279**, 7370–7377 (2004).

19. Hawlich, H. *et al.* C5a negatively regulates Toll-like receptor-4-induced immune responses. *Immunity* **22**, 415–426 (2005).
This study describes a negative effect of C5a on TLR4-mediated responses as an important mechanism for regulating T_H1-cell polarization in response to activation of the innate and adaptive immune systems.

20. Zhang, X. *et al.* Regulation of Toll-like receptor-mediated inflammatory response by complement *in vivo*. *Blood* **110**, 228–236 (2007).

21. Koleva, M. *et al.* Induction of anaphylatoxin C5a receptors in rat hepatocytes by lipopolysaccharide *in vivo*: mediation by interleukin-6 from Kupffer cells. *Gastroenterology* **122**, 697–708 (2002).

22. Clark, S. R. *et al.* Platelet TLR4 activates neutrophil extracellular traps to ensnare bacteria in septic blood. *Nature Med.* **13**, 463–469 (2007).

23. Daubeuf, B. *et al.* TLR4/MD-2 monoclonal antibody therapy affords protection in experimental models of septic shock. *J. Immunol.* **179**, 6107–6114 (2007).

24. McMasters, K. M., Peyton, J. C., Hadjiminis, D. J. & Cheadle, W. G. Endotoxin and tumour necrosis factor do not cause mortality from caecal ligation and puncture. *Cytokine* **6**, 530–536 (1994).

25. van der Poll, T. & Opal, S. M. Host–pathogen interactions in sepsis. *Lancet Infect. Dis.* **8**, 32–43 (2008).

26. Ward, P. A. The dark side of C5a in sepsis. *Nature Rev. Immunol.* **4**, 133–142 (2004).

27. Ritis, K. *et al.* A novel C5a receptor–tissue factor cross-talk in neutrophils links innate immunity to coagulation pathways. *J. Immunol.* **177**, 4794–4802 (2006).

28. Riedemann, N. C. *et al.* Regulatory role of C5a on macrophage migration inhibitory factor release from neutrophils. *J. Immunol.* **173**, 1355–1359 (2004).

29. Rittirsch, D. *et al.* Functional roles for C5a receptors in sepsis. *Nature Med.* **14**, 551–557 (2008).
This study indicates that both C5a receptors, C5AR and C5L2, synergistically contribute to harmful events in sepsis. In contrast to previous discussions, the authors conclude that C5L2 seems to be a functional receptor rather than a decoy receptor.

30. Calandra, T. & Roger, T. Macrophage migration inhibitory factor: a regulator of innate immunity. *Nature Rev. Immunol.* **3**, 791–800 (2003).

31. Lubetsky, J. B. *et al.* The tautomerase active site of macrophage migration inhibitory factor is a potential target for discovery of novel anti-inflammatory agents. *J. Biol. Chem.* **277**, 24976–24982 (2002).

32. Shi, X. *et al.* CD44 is the signaling component of the macrophage migration inhibitory factor–CD74 receptor complex. *Immunity* **25**, 595–606 (2006).

33. Bernhagen, J. *et al.* MIF is a pituitary-derived cytokine that potentiates lethal endotoxaemia. *Nature* **365**, 756–759 (1993).

34. Leng, L. *et al.* MIF signal transduction initiated by binding to CD74. *J. Exp. Med.* **197**, 1467–1476 (2003).

35. Mitchell, R. A., Metz, C. N., Peng, T. & Bucala, R. Sustained mitogen-activated protein kinase (MAPK) and cytoplasmic phospholipase A2 activation by macrophage migration inhibitory factor (MIF). Regulatory role in cell proliferation and glucocorticoid action. *J. Biol. Chem.* **274**, 18100–18106 (1999).

36. Calandra, T. *et al.* Protection from septic shock by neutralization of macrophage migration inhibitory factor. *Nature Med.* **6**, 164–170 (2000).
This study defines a crucial role for MIF in the pathogenesis of septic shock and identifies MIF as a new target for therapeutic intervention.

37. Al-Abed, Y. *et al.* ISO-1 binding to the tautomerase active site of MIF inhibits its pro-inflammatory activity and increases survival in severe sepsis. *J. Biol. Chem.* **280**, 36541–36544 (2005).

38. Muller, S. *et al.* New EMBO members' review. The double life of HMGB1 chromatin protein: architectural factor and extracellular signal. *EMBO J.* **20**, 4337–4340 (2001).

39. Wang, H. *et al.* HMGB-1 as a late mediator of endotoxin lethality in mice. *Science* **285**, 248–251 (1999).
This work defined, for the first time, an extracellular role for HMGB1 as a pro-inflammatory mediator in endotoxaemia and sepsis.

40. Lotze, M. T. & Tracey, K. J. High mobility group box 1 protein (HMGB1): nuclear weapon in the immune arsenal. *Nature Rev. Immunol.* **5**, 331–342 (2005).

41. Kim, J. Y. *et al.* HMGB1 contributes to the development of acute lung injury after hemorrhage. *Am. J. Physiol. Lung Cell. Mol. Physiol.* **288**, L958–L965 (2005).

42. Wang, H. *et al.* Cholinergic agonists inhibit HMGB1 release and improve survival in experimental sepsis. *Nature Med.* **10**, 1216–1221 (2004).

43. Scaffidi, P., Misteli, T. & Bianchi, M. E. Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. *Nature* **418**, 191–195 (2002).

44. Qin, S. *et al.* Role of HMGB1 in apoptosis-mediated sepsis lethality. *J. Exp. Med.* **203**, 1637–1642 (2006).

45. Hori, O. *et al.* The receptor for advanced glycation end products (RAGE) is a cellular binding site for amphotericin. Mediation of neurite outgrowth and co-expression of rage and amphotericin in the developing nervous system. *J. Biol. Chem.* **270**, 25752–25761 (1995).

46. Sha, Y., Zmijewski, J., Xu, Z. & Abraham, E. HMGB1 develops enhanced proinflammatory activity by binding to cytokines. *J. Immunol.* **180**, 2531–2537 (2008).

47. Klune, J. R., Dhupar, R., Cardinal, J., Billiar, T. R. & Tsung, A. HMGB1: endogenous danger signaling. *Mol. Med.* **14**, 476–484 (2008).

48. Sundén-Cullberg, J. *et al.* Persistent elevation of high mobility group box-1 protein (HMGB1) in patients with severe sepsis and septic shock. *Crit. Care Med.* **33**, 564–573 (2005).

49. Liliensiek, B. *et al.* Receptor for advanced glycation end products (RAGE) regulates sepsis but not the adaptive immune response. *J. Clin. Invest.* **113**, 1641–1650 (2004).

50. Yang, H. *et al.* Reversing established sepsis with antagonists of endogenous high-mobility group box 1. *Proc. Natl Acad. Sci. USA* **101**, 296–301 (2004).

51. Weaver, C. T., Hatton, R. D., Mangan, P. R. & Harrington, L. E. IL-17 family cytokines and the expanding diversity of effector T-cell lineages. *Annu. Rev. Immunol.* **25**, 821–852 (2007).

52. Flierl, M. A. *et al.* Adverse functions of IL-17A in experimental sepsis. *FASEB J.* **22**, 2198–2205 (2008).

53. Nakae, H. *et al.* Serum complement levels and severity of sepsis. *Res. Commun. Chem. Pathol. Pharmacol.* **84**, 189–195 (1994).

54. Gerard, C. Complement C5a in the sepsis syndrome — too much of a good thing? *N. Engl. J. Med.* **348**, 167–169 (2003).

55. Kildsgaard, J. *et al.* Cutting edge: targeted disruption of the C3a receptor gene demonstrates a novel protective anti-inflammatory role for C3a in endotoxin-shock. *J. Immunol.* **165**, 5406–5409 (2000).

56. Francis, K. *et al.* Complement C3a receptors in the pituitary gland: a novel pathway by which an innate immune molecule releases hormones involved in the control of inflammation. *FASEB J.* **17**, 2266–2268 (2003).

57. Huber-Lang, M. S. *et al.* Complement-induced impairment of innate immunity during sepsis. *J. Immunol.* **169**, 3223–3231 (2002).

58. Huber-Lang, M. *et al.* Role of C5a in multiorgan failure during sepsis. *J. Immunol.* **166**, 1193–1199 (2001).

59. Riedemann, N. C. *et al.* C5a receptor and thymocyte apoptosis in sepsis. *FASEB J.* **16**, 887–888 (2002).

60. Flierl, M. A. *et al.* The complement anaphylatoxin C5a induces apoptosis in adrenomedullary cells during experimental sepsis. *PLoS ONE* **3**, e2560 (2008).

61. Laudes, I. J. *et al.* Anti-c5a ameliorates coagulation/fibrinolytic protein changes in a rat model of sepsis. *Am. J. Pathol.* **160**, 1867–1875 (2002).

62. Niederbichler, A. D. *et al.* An essential role for complement C5a in the pathogenesis of septic cardiac dysfunction. *J. Exp. Med.* **203**, 53–61 (2006).

63. Gerard, N. P. *et al.* An anti-inflammatory function for the complement anaphylatoxin C5a-binding protein, C5L2. *J. Biol. Chem.* **280**, 39677–39680 (2005).

64. Chen, N. J. *et al.* C5L2 is critical for the biological activities of the anaphylatoxins C5a and C3a. *Nature* **446**, 203–207 (2007).

65. Huber-Lang, M. *et al.* Changes in the novel orphan, C5a receptor (C5L2), during experimental sepsis and sepsis in humans. *J. Immunol.* **174**, 1104–1110 (2005).

66. Zeerleder, S. *et al.* Administration of C1 inhibitor reduces neutrophil activation in patients with sepsis. *Clin. Diagn. Lab. Immunol.* **10**, 529–535 (2003).

67. Caliezi, C. *et al.* C1-inhibitor in patients with severe sepsis and septic shock: beneficial effect on renal dysfunction. *Crit. Care Med.* **30**, 1722–1728 (2002).

68. Flierl, M. A. *et al.* Functions of the complement components C3 and C5 during sepsis. *FASEB J.* 27 June 2008 (doi:10.1096/fj.08-110595).

69. Levi, M. & Ten Cate, H. Disseminated intravascular coagulation. *N. Engl. J. Med.* **341**, 586–592 (1999).

70. Abraham, E. Coagulation abnormalities in acute lung injury and sepsis. *Am. J. Respir. Cell. Mol. Biol.* **22**, 401–404 (2000).

71. Esmon, C. T. The impact of the inflammatory response on coagulation. *Thromb. Res.* **114**, 321–327 (2004).

72. Stouthard, J. M. *et al.* Interleukin-6 stimulates coagulation, not fibrinolysis, in humans. *Thromb. Haemost.* **76**, 738–742 (1996).

73. Bevilacqua, M. P. *et al.* Recombinant tumor necrosis factor induces procoagulant activity in cultured human vascular endothelium: characterization and comparison with the actions of interleukin-1. *Proc. Natl Acad. Sci. USA* **83**, 4533–4537 (1986).

74. Bernard, G. R. *et al.* Efficacy and safety of recombinant human activated protein C for severe sepsis. *N. Engl. J. Med.* **344**, 699–709 (2001).
This study showed that treatment with activated protein C significantly decreased mortality in patients with severe sepsis, but might be associated with an increased risk of bleeding.

75. Maugeri, N. *et al.* Human polymorphonuclear leukocytes produce and express functional tissue factor upon stimulation. *J. Thromb. Haemost.* **4**, 1323–1330 (2006).

76. Osterud, B. & Flaegstad, T. Increased tissue thromboplastin activity in monocytes of patients with meningococcal infection: related to an unfavourable prognosis. *Thromb. Haemost.* **49**, 5–7 (1983).

77. Zeerleder, S., Schroeder, V., Hack, C. E., Kohler, H. P. & Willemin, W. A. TAFI and PAI-1 levels in human sepsis. *Thromb. Res.* **118**, 205–212 (2006).

78. Levi, M., de Jonge, E. & van der Poll, T. New treatment strategies for disseminated intravascular coagulation based on current understanding of the pathophysiology. *Ann. Med.* **36**, 41–49 (2004).

79. Esmon, C. T. The protein C pathway. *Chest* **124**, 265–325 (2003).

80. Joyce, D. E., Gelbert, L., Ciaccia, A., DeHoff, B. & Grinnell, B. W. Gene expression profile of antithrombotic protein C defines new mechanisms modulating inflammation and apoptosis. *J. Biol. Chem.* **276**, 11199–11203 (2001).

81. Riewald, M., Petrovan, R. J., Donner, A., Mueller, B. M. & Ruf, W. Activation of endothelial cell protease activated receptor 1 by the protein C pathway. *Science* **296**, 1880–1882 (2002).

82. Niessen, F. *et al.* Dendritic cell PAR1–S1P3 signalling couples coagulation and inflammation. *Nature* **452**, 654–658 (2008).

83. Kaneider, N. C. *et al.* 'Role reversal' for the receptor PAR1 in sepsis-induced vascular damage. *Nature Immunol.* **8**, 1303–1312 (2007).

84. Fukudome, K. & Esmon, C. T. Identification, cloning, and regulation of a novel endothelial cell protein C/activated protein C receptor. *J. Biol. Chem.* **269**, 26486–26491 (1994).

85. Ito, T. *et al.* High-mobility group box 1 protein promotes development of microvascular thrombosis in rats. *J. Thromb. Haemost.* **5**, 109–116 (2007).

86. Cheng, T. *et al.* Activated protein C blocks p53-mediated apoptosis in ischemic human brain endothelium and is neuroprotective. *Nature Med.* **9**, 338–342 (2003).

87. Kerschen, E. J. *et al.* Endotoxemia and sepsis mortality reduction by non-anticoagulant activated protein C. *J. Exp. Med.* **204**, 2439–2448 (2007).

88. Krem, M. M. & Di Cera, E. Evolution of enzyme cascades from embryonic development to blood coagulation. *Trends Biochem. Sci.* **27**, 67–74 (2002).

89. Ghebrehiwet, B., Silverberg, M. & Kaplan, A. P. Activation of the classical pathway of complement by Hageman factor fragment. *J. Exp. Med.* **153**, 665–676 (1981).

90. Goldberger, G. *et al.* N_H2-terminal structure and cleavage of guinea pig pro-C3, the precursor of the third complement component. *J. Biol. Chem.* **256**, 12617–12619 (1981).

91. Thoman, M. L., Meuth, J. L., Morgan, E. L., Weigle, W. O. & Hugli, T. E. C3d-K, a kallikrein cleavage fragment of iC3b is a potent inhibitor of cellular proliferation. *J. Immunol.* **133**, 2629–2633 (1984).

92. Campbell, W., Okada, N. & Okada, H. Carboxypeptidase R is an inactivator of complement-derived inflammatory peptides and an inhibitor of fibrinolysis. *Immunol. Rev.* **180**, 162–167 (2001).
93. Wojta, J. *et al.* C5a stimulates production of plasminogen activator inhibitor-1 in human mast cells and basophils. *Blood* **100**, 517–523 (2002).
94. Muhlfelder, T. W. *et al.* C5 chemotactic fragment induces leukocyte production of tissue factor activity: a link between complement and coagulation. *J. Clin. Invest.* **63**, 147–150 (1979).
95. Krarup, A., Wallis, R., Presanis, J. S., Gal, P. & Sim, R. B. Simultaneous activation of complement and coagulation by MBL-associated serine protease 2. *PLoS ONE* **2**, e623 (2007).
96. Rezende, S. M., Simmonds, R. E. & Lane, D. A. Coagulation, inflammation, and apoptosis: different roles for protein S and the protein S–C4b binding protein complex. *Blood* **103**, 1192–1201 (2004).
97. Sternberg, E. M. Neural regulation of innate immunity: a coordinated nonspecific host response to pathogens. *Nature Rev. Immunol.* **6**, 318–328 (2006).
98. Elenkov, I. J., Wilder, R. L., Chrousos, G. P. & Vizi, E. S. The sympathetic nerve — an integrative interface between two supersystems: the brain and the immune system. *Pharmacol. Rev.* **52**, 595–638 (2000).
99. Bergquist, J., Tarkowski, A., Ekman, R. & Ewing, A. Discovery of endogenous catecholamines in lymphocytes and evidence for catecholamine regulation of lymphocyte function via an autocrine loop. *Proc. Natl Acad. Sci. USA* **91**, 12912–12916 (1994).
100. Munford, R. S. & Tracey, K. J. Is severe sepsis a neuroendocrine disease? *Mol. Med.* **8**, 437–442 (2002).
101. Wang, H. *et al.* Nicotinic acetylcholine receptor $\alpha 7$ subunit is an essential regulator of inflammation. *Nature* **421**, 384–388 (2003).
This work shows that the α_7 -nicotinic acetylcholine receptor is required for acetylcholine-mediated inhibition of cytokine production by macrophages, which is also known as the cholinergic anti-inflammatory pathway.
102. Tracey, K. J. Physiology and immunology of the cholinergic anti-inflammatory pathway. *J. Clin. Invest.* **117**, 289–296 (2007).
103. Huston, J. M. *et al.* Splenectomy inactivates the cholinergic anti-inflammatory pathway during lethal endotoxemia and polymicrobial sepsis. *J. Exp. Med.* **203**, 1623–1628 (2006).
104. Huston, J. M. *et al.* Transcutaneous vagus nerve stimulation reduces serum high mobility group box 1 levels and improves survival in murine sepsis. *Crit. Care Med.* **35**, 2762–2768 (2007).
105. Annane, D. *et al.* Inappropriate sympathetic activation at onset of septic shock: a spectral analysis approach. *Am. J. Respir. Crit. Care Med.* **160**, 458–465 (1999).
106. Bergmann, M. & Sautner, T. Immunomodulatory effects of vasoactive catecholamines. *Wien. Klin. Wochenschr.* **114**, 752–761 (2002).
107. Oberbeck, R. Catecholamines: physiological immunomodulators during health and illness. *Curr. Med. Chem.* **13**, 1979–1989 (2006).
108. Oberbeck, R. *et al.* Adrenergic modulation of survival and cellular immune functions during polymicrobial sepsis. *Neuroimmunomodulation* **11**, 214–223 (2004).
109. Kradin, R., Rodberg, G., Zhao, L. H. & Leary, C. Epinephrine yields translocation of lymphocytes to the lung. *Exp. Mol. Pathol.* **70**, 1–6 (2001).
110. Ackerman, K. D., Madden, K. S., Livnat, S., Felten, S. Y. & Felten, D. L. Neonatal sympathetic denervation alters the development of *in vitro* spleen cell proliferation and differentiation. *Brain Behav. Immun.* **5**, 235–261 (1991).
111. Spengler, R. N., Allen, R. M., Remick, D. G., Strieter, R. M. & Kunkel, S. L. Stimulation of α -adrenergic receptor augments the production of macrophage-derived tumor necrosis factor. *J. Immunol.* **145**, 1430–1434 (1990).
112. Freestone, P. P. *et al.* Growth stimulation of intestinal commensal *Escherichia coli* by catecholamines: a possible contributory factor in trauma-induced sepsis. *Shock* **18**, 465–470 (2002).
113. Zhou, M., Das, P., Simms, H. H. & Wang, P. Gut-derived norepinephrine plays an important role in up-regulating IL-1 β and IL-10. *Biochim. Biophys. Acta* **1740**, 446–452 (2005).
114. Yang, S., Zhou, M., Chaudry, I. H. & Wang, P. Norepinephrine-induced hepatocellular dysfunction in early sepsis is mediated by activation of α_2 -adrenoceptors. *Am. J. Physiol. Gastrointest. Liver Physiol.* **281**, G1014–G1021 (2001).
115. Yang, S., Koo, D. J., Zhou, M., Chaudry, I. H. & Wang, P. Gut-derived norepinephrine plays a critical role in producing hepatocellular dysfunction during early sepsis. *Am. J. Physiol. Gastrointest. Liver Physiol.* **279**, G1274–G1281 (2000).
116. Zhou, M. *et al.* The role of Kupffer cell α_2 -adrenoceptors in norepinephrine-induced TNF- α production. *Biochim. Biophys. Acta* **1537**, 49–57 (2001).
117. Dombrovskiy, V. Y., Martin, A. A., Sunderram, J. & Paz, H. L. Rapid increase in hospitalization and mortality rates for severe sepsis in the United States: a trend analysis from 1993 to 2003. *Crit. Care Med.* **35**, 1244–1250 (2007).
This publication is the most recent study to describe the epidemiology of sepsis in the United States. Most importantly, the rates of hospitalization and mortality from severe sepsis increased significantly over the observation period from 1993 to 2003.
118. Rittirsch, D., Hoesel, L. M. & Ward, P. A. The disconnect between animal models of sepsis and human sepsis. *J. Leukocyte Biol.* **81**, 137–143 (2007).
119. Levy, M. M. *et al.* 2001 SCCM/ESICM/ACCP/ATS/SIS International Sepsis Definitions Conference. *Crit. Care Med.* **31**, 1250–1256 (2003).
120. Calvano, S. E. *et al.* A network-based analysis of systemic inflammation in humans. *Nature* **437**, 1032–1037 (2005).
121. Rossaint, R. *et al.* Key issues in advanced bleeding care in trauma. *Shock* **26**, 322–331 (2006).
122. Westfall, T. C. & Westfall, D. P. in *Goodman & Gilman's The Pharmacological Basis of Therapeutics* (eds Brunton, L. L., Lazo, J. S. & Parker, K. L.) 153–158 (McGraw-Hill, New York, 2006).

Acknowledgements

This work was supported by grants GM-29,507, HL-31963 and GM-61656 from the National Institutes of Health, USA, to P.A.W.

DATABASES

Entrez Gene: <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=gene>
antithrombin III | C3AR | C5AR | C5L2 | CD14 | CD44 | CD74 | EPCR | HMGB1 | IFN γ | IL-1 β | IL-4 | IL-5 | IL-6 | IL-10 | IL-12 | IL-13 | IL-17A | MASP2 | MD2 | MIF | PAI1 | PAR1 | protein C | protein S | RAGE | TAEI | TEPI | thrombin | tissue factor | TLR2 | TLR4 | TNF | TPA

FURTHER INFORMATION

Peter Ward's homepage: <http://www.med.umich.edu/immprog/faculty/wardp.htm>

ALL LINKS ARE ACTIVE IN THE ONLINE PDF